



# Revista Colombiana de Anestesiología

## Colombian Journal of Anesthesiology

www.revcolanest.com.co



## Questions and answers

### Fernando Raffán Sanabria\*

Anestesiólogo intensivista, Hospital Universitario Fundación Santa Fe de Bogotá, Colombia  
 Profesor, Universidad el Bosque y Universidad de los Andes, Bogotá, Colombia

This section includes questions prepared on the basis of the contents of the articles published in volume 40, number 3 of the *Colombian Journal of Anesthesiology*. Accept the challenge to test your reading comprehension and knowledge.

#### Instructions

This is a multiple-choice questionnaire. Please mark the appropriate answer.

- 1) The most frequent cause of lumbar facet syndrome is related to:
  - A) Disc hernia
  - B) Lumbar canal stenosis
  - C) Radiculopathy
  - D) Degenerative joint disease
- 2) The most frequent sensitive syndrome of the lumbar facet syndrome is:
  - A) Lower limb paresthesia
  - B) Lower limb dysesthesia
  - C) A combination of aresthesia and dysisthesia
  - D) Radiating pain
- 3) The following characteristics are present in the so called Tokobsubo cardiomyopathy except for:
  - A) More frequent in females
  - B) Frequently exhibits ST segment elevation in the ECG
  - C) Usually associated with coronary disease
  - D) Elevation of enzymatic markers
- 4) The following drugs have been associated with ST segment elevation, except for:
  - A) Propofol
  - B) Nitroglycerin
  - C) Etomidate
  - D) Oxitocin
- 5) The so called facet block for treating the facet syndrome consists in the administration of:
  - A) A steroid
  - B) A local anesthetic
  - C) A 2-drug combination
  - D) Absolute alcohol plus steroid administered together
- 6) The recommended dose volume for the application of a solution for facet block should not exceed:
  - A) 1 c.c.
  - B) 2 c.c..
  - C) 3 c.c..
  - D) 4 c.c.
- 7) Which of the following mechanisms has been attributed to the genesis of coronary spasm:
  - A) Decrease of serotonin endogenous ligands
  - B) Stimulation of the nitric oxide synthase enzyme
  - C) Activation of the K ATP channels
  - D) Calcium channels genetic mutation
- 8) Among the strategies to prevent a new intraoperative coronary spasm event, the following are included, except for:
  - A) IV calcium administration
  - B) IV magnesium
  - C) ACE inhibitors
  - D) Clopidogrel

\* Corresponding author at: Calle119 # 7 - 75, Bogotá, Colombia.  
 E-mail address: raffanmago@gmail.com

- 9) Cesarean delivery morbidity and mortality studies under general anesthesia (GA) vs. regional anesthesia (RA) have shown that:
- A) Maternal mortality is significantly higher with GA than with RA
  - B) The percentage of patients who ambulate before 24 h following C-section is higher with GA than with RA.
  - C) The umbilical vein pH is lower with GA versus RA
  - D) The neurological adaptive ability of the newborn is significantly lower with GA versus RA
- 10) The Food and Drug Administration (FDA) of the United States recommends classifying drugs during pregnancy in terms of a risk: benefit determination. Such classification establishes that the following drug falls in the X category
- A) Diazepam
  - B) Diclofenac
  - C) Naloxone
  - D) Ergot byproducts
- 11) In terms of the new cell model of coagulation, the following statements are true:
- A) The interaction between Factor VIII and the tissue factor is the key process in the initiation phase
  - B) During the amplification phase the negative feedback system is activated through natural anticoagulants such as protein C and antithrombin
  - C) During the propagation phase, the prothrombin complex is inhibited due to factor X activation
  - D) Platelet activation increases the permeability of the membrane as a result of Na releasing Factor VII and phospholipids
- 12) Which of the following parameters account for a higher thrombin output in the thromboelastogram:
- A) R interval
  - B) K interval
  - C) Alpha angle
  - D) Maximum amplitude

---

## Reference

Raffán F. Preguntas y respuestas. Rev. colomb. anestesiología. 2012;40(3).

### Answers:

- 1) D
- 2) A
- 3) C
- 4) C
- 5) C
- 6) A
- 7) C
- 8) A
- 9) C
- 10) D
- 11) B
- 12) A